

Mold Avoiders vs. Other Other Groups

Mold Avoiders: n = 66

Other groups: n = 92

Participants from the Mold Avoiders group were significantly more likely to say that they had benefited from the following therapies than were participating members of the other groups.

	<u>Mold Avoiders</u>	<u>Other Groups</u>	<u>P-value</u>
Avoiding toxic buildings	92%	74%	p<.01
Avoiding cross-contaminated possessions	74%	49%	p<.01
Avoiding toxic locations	63%	47%	p<.05
Rapid decontamination (e.g. (within minutes of exposure)	21%	2%	p<.001
Consistent decontamination before returning to safe space	21%	10%	p<.10
Avoiding glyphosate in food	27%	11%	p<.01
Fermented foods	23%	7%	p<.01
Methylation related supplements	38%	14%	p<.001
Ionic detox footbath	17%	4%	p<.01
Homeopathy	24%	11%	p<.05
Neural therapy	8%	1%	p<.05

P-values were calculated using a Z-test.

Participants from the Mold Avoiders group were significantly more likely to say that they had benefited from the following therapies than were participating members of the other groups.

	<u>Mold Avoiders</u>	<u>Other Groups</u>	<u>P-value</u>
Reducing stress	27%	48%	p<.01
Probiotics	8%	43%	p<.001
Activated charcoal	14%	39%	p<.001
Glutathione	15%	30%	p<.05
Avoiding foods with mycotoxins and molds	9%	30%	p<.01
Antifungals	3%	21%	p<.01